WHAT IS CLAIMED IS:

Polymer 1

1. A composition of matter comprising a polymer of the general structure:

CH₃O CH₃O CH₃O OCH₃

C/R2/D

A/R1/B

E/R3/F

-(CH₂)_x-/-O(CH₂)_x -/-O(CH₂)_xO-

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para with respect to the pyridyl nitrogen;

wherein bonds C and D may be either ortho, meta or para with respect one another; and

wherein bonds E and F may be either ortho, meta or para with respect one another;

wherein Y may be a moiety selected from the group consisting of $-(CH_2)_x$ - $-(CH_2)_x$ O-, $-O(CH_2)_x$ – and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive; and

wherein n is an integer greater than 1.

35

10

- 2. A composition according to claim 1 wherein at least one R2 substituent is a methoxy group.
- 5 3. A composition according to claim 1 wherein at least two R2 substituents are methoxy groups.
 - 4. A composition according to claim 1 wherein at least one R3 substituent is a methoxy group.
 - 5. A composition according to claim 1 wherein at least two R3 substituents are methoxy groups.
- 6. A composition according to claim 1 wherein vinyl linkage A attaches at a position ortho to the pyridyl nitrogen.
 - 7. A composition according to claim 1 wherein vinyl linkage B attaches at a position ortho to the pyridyl nitrogen.
 - 8. A composition according to claim 1 wherein vinyl linkage A attaches at a position para to the pyridyl nitrogen.
 - 9. A composition according to claim 1 wherein vinyl linkage B attaches at a position para to the pyridyl nitrogen.
 - 10. A composition according to claim 1 wherein x is an integer in the range of 1 to 6 inclusive.

Polymer 2

11. A composition of matter comprising a polymer of the general structure:

R4 C/R2/D - A/R1/B E/R3/F

 $-(CH_2)_x$ -/- $O(CH_2)_x$ -/- $O(CH_2)_x$ O-

wherein

5

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may be either ortho or para with respect to the pyridyl nitrogen;

wherein bonds C and D may independently be either ortho, meta or para with respect one another;

wherein bonds E and F may be either ortho, meta or para with respect one another;

wherein bonds G and H may be either ortho, meta or para with respect one another;

wherein Y may be a moiety selected from the group consisting of $-(CH_2)_x$, $-(CH_2)_x$ O-, $-O(CH_2)_x$ – and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive;

wherein Z may be a moiety selected from the group consisting of $-(CH_2)_x$, $-(CH_2)_x$ O-, $-O(CH_2)_x$ — and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive; and

- wherein n is an integer greater than 1.
 - 12. A composition according to claim 11 wherein at least one R2 substituent is a methoxy group.
- 40 13. A composition according to claim 11 wherein at least two R2 substituents are methoxy groups.
 - 14. A composition according to claim 11 wherein at least one R3 substituent is a methoxy group.

- 15. A composition according to claim 11 wherein at least two R3 substituents are methoxy groups.
- 16. A composition according to claim 11 wherein at least one R3 substituent is a methoxy group.
 - 17. A composition according to claim 11 wherein at least two R3 substituents are methoxy groups.
- 10 18. A composition according to claim 11 wherein vinyl linkage A attaches at a position ortho to the pyridyl nitrogen.
 - 19. A composition according to claim 11 wherein vinyl linkage B attaches at a position ortho to the pyridyl nitrogen.
 - 20. A composition according to claim 11 wherein vinyl linkage A attaches at a position para to the pyridyl nitrogen.
 - 21. A composition according to claim 11 wherein vinyl linkage B attaches at a position para to the pyridyl nitrogen.
 - 22. A composition according to claim 11 wherein x is an integer in the range of 1 to 6 inclusive.

Oligomers 1, 2, 3 & 4

23. A composition of matter comprising an oligomer of the general structure:

35 wherein

15

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;


wherein bond C may be either ortho, meta or para with respect to the respective quinoyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective quinoyl nitrogen.

- 24. A composition according to claim 23 wherein at least one R1 substituent is a methoxy group.
- 25. A composition according to claim 23 wherein at least two R1 substituents are methoxy groups.

Oligomer 5, 6, 7 & 8

26. A composition of matter comprising an oligomer of the general structure:



R2 C A R1 B D R3

wherein

5

Sand the family from

30

35


- the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- wherein bonds A and B may independently be either ortho, meta or para from one another;
 - wherein bond C may be either ortho, meta or para with respect to the respective pyridyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective pyridyl nitrogen.

- 27. A composition according to claim 26 wherein at least one R1 substituent is a methoxy group.
- 28. A composition according to claim 26 wherein at least two R1 substituents are methoxy groups.
- 29. A composition according to claim 26 wherein at least one R2 substituent is a methyl group.
- 30. A composition according to claim 26 wherein at least one R3 substituent is a methyl group.

Oligomers 9 & 10

31. A composition of matter comprising an oligomer of the general structure:



R3/R2 C D R1 R4/R5

5

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

10

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of

unite tand the limb that the land that the land the

20

T125

hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of

hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may be either ortho, meta or para from one another.

32. A composition according to claim 31 wherein at least one R1 substituent is a methoxy group.


33. A composition according to claim 31 wherein at least two R1 substituents are

methoxy groups.

30

Oligomers 11, 12 & 13

A composition of matter comprising an oligomer of the general structure: 34.



wherein


- the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups; and

wherein bonds A and B may be either ortho or para from the pyridyl nitrogen.

- 35. A composition according to claim 34 wherein at least one R2 substituent is a methoxy group.
- 36. A composition according to claim 34 wherein two R2 substituents are methoxy groups.
- 37. A composition according to claim 34 wherein three R2 substituents are methoxy groups.
- 38. A composition according to claim 34 wherein at least one R3 substituent is a methoxy group.
- 39. A composition according to claim 34 wherein two R3 substituents are methoxy groups.
- 40. A composition according to claim 34 wherein three R2 substituents are methoxy groups.

Block Co-polymer of Oligomers 1, 2, 3 & 4 (Y only)

35 41. A composition of matter comprising a block co-polymer of the general structure:



wherein

5

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;

wherein bond C may be either ortho, meta or para with respect to the respective quinoyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective quinoyl nitrogen;

wherein Y may be a moiety attached at any point on rings R2 and R3, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-$, $-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive; and

wherein n is an integer greater than 1.

35

25

42. A composition according to claim 41 wherein at least one R1 substituent is a methoxy group.

30

10

- 43. A composition according to claim 41 wherein at least two R1 substituents are methoxy groups.
- wherein Y may be a moiety selected from the group consisting of $-(CH_2)_x$ -, $-(CH_2)_x$ O-, $-O(CH_2)_x$ and $-O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive;

Block Co-polymer of Oligomers 1, 2, 3 & 4 (Y, R & Z)

44. A composition of matter comprising a block co-polymer of the general structure:

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R6 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;

35

10

15

wherein bond C may be either ortho, meta or para with respect to the respective quinoyl nitrogen; and

wherein bond D may be either ortho, meta or para with respect to the respective quinoyl nitrogen;

wherein Y may be a moiety attached at any point on ring R6, and may be selected from the group consisting of $-(CH_2)_x--(CH_2)_xO-$, $-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;

wherein Z may be a moiety bridging any two points on rings R2 or R3 and R6, and may be selected from the group consisting of $-(CH_2)_{x}$, $-(CH_2)_{x}$ O-, $-O(CH_2)_{x}$ – and – $O(CH_2)_{x}$ O- wherein x is an integer in the range of 1 to 15 inclusive; and

wherein n is an integer greater than 1.

- 45. A composition according to claim 44 wherein at least one R1 substituent is a methoxy group.
- 46. A composition according to claim 44 wherein at least two R1 substituents are methoxy groups.

Block Co-polymer of Oligomers 5, 6, 7 & 8 (Y only)

47. A composition of matter comprising a block co-polymer of the general structure:

wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

15

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;

wherein bond C may be either ortho, meta or para with respect to the respective pyridyl nitrogen;

wherein bond D may be either ortho, meta or para with respect to the respective pyridyl nitrogen;

wherein Y may be a moiety attached at any point on ring R2, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-,-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;

and

wherein n is an integer greater than 1.

- 48. A composition according to claim 47 wherein at least one R1 substituent is a methoxy group.
- 49. A composition according to claim 47 wherein at least two R1 substituents are methoxy groups.
- 50. A composition according to claim 47 wherein at least one R2 substituent is a methyl group.
- 51. A composition according to claim 47 wherein at least one R3 substituent is a methyl group.

Block Co-polymer of Oligomers 5, 6, 7 & 8 (Y, R & Z)


52. A composition of matter comprising a block co-polymer of the general structure:

20

25

30

35



R2 C A R1 B D R3

5 wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may independently be either ortho, meta or para from one another;

wherein bond C may be either ortho, meta or para with respect to the respective pyridyl nitrogen;

wherein bond D may be either ortho, meta or para with respect to the respective pyridyl nitrogen;

wherein Y may be a moiety attached at any point on ring R4, and may be selected from the group consisting of $-(CH_2)_x--(CH_2)_xO-$, $-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;

wherein Z may be a moiety bridging any two points on rings R2 and R4, and may be selected from the group consisting of $-(CH_2)_x$ -, $-(CH_2)_x$ O-, $-O(CH_2)_x$ – and - $O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive; and

and

wherein n is an integer greater than 1.


40 53. A composition according to claim 52 wherein at least one R1 substituent is a methoxy group.

10

- 54. A composition according to claim 52 wherein at least two R1 substituents are methoxy groups.
- 5 55. A composition according to claim 52 wherein at least one R2 substituent is a methyl group.
 - 56. A composition according to claim 52 wherein at least one R3 substituent is a methyl group.

Block Co-polymer of Oligomers 9 & 10 (Y only)

57. A composition of matter comprising a block co-polymer of the general structure:



wherein

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

- 25 the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

wherein bonds A and B may be either ortho, meta or para from one another;


wherein Y may be a moiety attached at any point on rings R2 or R3, and may be selected from the group consisting of $-(CH_2)_x$, $-(CH_2)_x$ O-, $-O(CH_2)_x$ — and - $O(CH_2)_x$ O- wherein x is an integer in the range of 1 to 15 inclusive;

and

- wherein n is an integer greater than 1.
 - 58. A composition according to claim 57 wherein at least one R1 substituent is a methoxy group.
- 15 59. A composition according to claim 57 wherein at least two R1 substituents are methoxy groups.

Block Co-polymer of Oligomers 9 & 10 (Y, R and Z only)

60. A composition of matter comprising a block co-polymer of the general structure:



R3/R2 C A R1 B D R4/R5

25 wherein

30

the R1 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

- the R2 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R3 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;

15

- the R4 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
- the R5 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - the R6 substituents are independently selected from the group consisting of hydrogen, alkyl groups, alkoxy groups, and aryl groups;
 - wherein bonds A and B may be either ortho, meta or para from one another;
 - wherein Y may be a moiety attached at any point on ring R6, and may be selected from the group consisting of $-(CH_2)_x-,-(CH_2)_xO-,-O(CH_2)_x-$ and $-O(CH_2)_xO-$ wherein x is an integer in the range of 1 to 15 inclusive;
 - wherein Z may be a moiety bridging any two points on rings R2 or R3 and R6, and may be selected from the group consisting of $-(CH_2)_{x^-}, -(CH_2)_{x^-}, -(CH_2)_{x^-}, -(CH_2)_{x^-}$ and $-(CH_2)_{x^-}, -(CH_2)_{x^-}$

and

wherein n is an integer greater than 1.